(Thayetmyo-Minbu border, 5,000 feet, Southern Chin Hills, Burma), not *Tribura thoracica saturata* Yen, Ornith. Monatsber., 1933, p. 16 (Yao-shan, Kwangsi), I propose

## Bradypterus luteoventris ticehursti, nom. nov.

Delacour has noted (Ibis, 1943, p. 35) that Ticehurst's name cannot be used but, doubting the validity of the form, has neglected to re-name it. A single winter specimen from the mountains of northern Siam, however, is distinct from both *luteoventris* and *russulus* and agrees perfectly with Ticehurst's description of this putative race.

H. G. DEIGNAN.

## HOYT'S HORNED LARK ON THE EASTERN SHORE, MARYLAND.4

An adult male of Eremophila (Chionophilos) alpestris hoyti was collected at Cambridge, Dorchester county, Maryland, February 4, 1918, by Ralph W. Jackson, who presented it to the United States National Museum in 1920, together with an example of Eremophila a. alpestris of the same place and date. The former is now registered as U. S. N. M. no. 256939. The specimen has at some time been identified and initialed by H. C. Oberholser but seems never to have been noticed in print. The race is not otherwise recorded from Maryland and has apparently not previously been found farther south than central Ohio and Long Island.

H. G. DEIGNAN.

## THE EUROPEAN WIDGEON NEAR WASHINGTON.

In the Proceedings of the Biological Society of Washington for March 25, 1929, May T. Cooke, in her paper "Birds of the Washington, D. C. region," lists three records for *Mareca penelope* from the Washington, D. C. area. To these I would add a fourth, which, while, not coming precisely within the generally accepted confines, is yet near enough (forty miles distant) I believe to be recorded.

It is an immature male just coming into its full plumage. It was taken by myself on Nanjemoy Creek, Charles Co., Maryland, November 3, 1941.

E. M. HASBROUCK.

<sup>4</sup> Published with permission of the Secretary of the Smithsonian Institution.

